

AN INVENTORY OF ZOOXANTHELLATE SCLERACTINIAN CORALS IN SINGAPORE, INCLUDING 33 NEW RECORDS

Danwei Huang

*Department of Biological Sciences, National University of Singapore,
14 Science Drive 4, Singapore 117543, Republic of Singapore
Scripps Institution of Oceanography, University of California, San Diego, 9500 Gilman Drive, La Jolla,
CA 92093-0208, U.S.A.
Email: huangdanwei@nus.edu.sg*

Karenne P. P. Tun, L. M. Chou and Peter A. Todd

*Department of Biological Sciences, National University of Singapore,
14 Science Drive 4, Singapore 117543, Republic of Singapore
Email: karenne@nus.edu.sg (KPPT); dbsclm@nus.edu.sg (LMC); dbspat@nus.edu.sg (PAT).*

ABSTRACT. – Records of Scleractinia in Singapore have not been updated for 14 years. We present an inventory of zooxanthellate hard coral species by consolidating past work, coral collection records and recent publications, as well as conducting field surveys at eight coral reefs south of mainland Singapore. Species assessment surveys and recent literature revealed a total of 161 species. Using updated taxonomy, and including 33 new records, we report an increase in the number of species found in Singapore from 187 to 255. Raffles Lighthouse registered the largest number of species and represents the most undisturbed reefs of the sites studied. No clear relationship can be established between species richness and distance from the Mainland. The total number of species in Singapore is comparable to the reefs in neighboring countries if reef area is taken into account. Only 63.1% of total species recorded have been found in recent years, but this is not an exhaustive survey. As 12.9% of all species have been discovered in the last four years, and only 52.4% of species with distribution ranges encompassing Singapore have been found, more new records can be expected from further assessments.

KEY WORDS. – Coral reef, distribution, Scleractinia, Singapore, species list, zooxanthellae.

INTRODUCTION

Singapore is situated in the vicinity of the Indo-West Pacific center of marine biodiversity, which is home to the highest number of scleractinian coral species in the world (Ekman, 1953; Briggs, 1999, 2000; Veron, 2000; Spalding et al., 2001; Carpenter & Springer, 2005; Barber et al., 2006). The Philippines and Indonesia, for instance, host more than 400 species each, while 348 species have been recorded off the coasts of Malaysia (Spalding et al., 2001). Yet in Singapore, flanked by the reefs of Malaysia and Indonesia, only 197 hard coral species have been identified in the last assessments concluded 14 years ago (Chua & Chou, 1991; Chou et al., 1994b; Chou et al., 1994c; Loo & Chou, 1995). This count was based on line intercept transects at specific depths of the reef (Dartnall & Jones, 1986), but studies of this nature mainly address questions regarding coral abundance (English et al., 1994). As a result, rare species are likely to be omitted from such surveys, rendering this technique inappropriate for the appraisal of species richness. Since

new scleractinian species have been described in recent years (e.g. Veron, 2002), the previous record of 197 species may be a considerable underestimation of the actual alpha diversity.

Coral reefs in Singapore have been impacted by the high rates of sedimentation resulting from land reclamation activities that commenced in 1963 (Chou, 1988a; Chou, 1996; Chou, 2006). Sediment deposition, suspended particles and associated light attenuation adversely affect the reproduction, recruitment, growth and survival of reef-building zooxanthellate (or hermatypic) corals (Roy & Smith, 1971; Endean, 1976; Rogers, 1990; Titlyanov & Latypov, 1991; Gilmour, 2002; Dikou & van Woesik, 2006; but see Lane, 1991). Singapore reefs tend not to thrive deeper than 3 m below the crest level, and reef accretion is thus limited to the shallows (Chou, 1988b; Goh & Chou, 1993). As coastal development continues unabated, there is a pressing need to assess and manage the hermatypic species that surround the offshore islands.

Records of Singapore's reef corals date back to the 1960s, when the first comprehensive list was published by Chuang (1961). It documented 63 zooxanthellate scleractinian species found along the Peninsula Malaysia coastline, of which Singapore's mainland and offshore islands accounted for the longest natural shore length prior to the systematic land reclamation (Chuang, 1961). Voucher specimens have also been collected during the intervening 48 years and deposited at the Zoological Reference Collection (ZRC), Raffles Museum of Biodiversity Research (RMBR), National University of Singapore (NUS). Taxonomic work by Koh & Chou (1989) and Leow (1992) revealed 13 species of Fungiidae and 18 species of *Acropora* respectively. Unfortunately, the inventory of 187 zooxanthellate species produced by Loo & Chou (1995) (ten less than the 197 hard coral records: three azooxanthellate, three non-scleractinian and four misnamed) did not account for the above. We therefore examined all relevant literature, revalidated museum records, and conducted surveys at selected reef sites using a more suitable method for species assessment. The primary objective of this study is to update the species inventory of zooxanthellate coral species in Singapore and present the new species records obtained since 2005.

MATERIAL AND METHODS

Field surveys. – Surveys were conducted between July 2006 and July 2007 at eight coral reefs fringing the southern offshore islands of Singapore: (1) Raffles Lighthouse (or Pulau [= Island] Satumu; two sites; 01°09'36.5''N 103°44'24.0''E; 01°09'37.0''N 103°44'29.0''E); (2) Pulau Semakau (01°12'11.5''N 103°45'15.0''E); (3) Pulau Jong (01°12'53.5''N 103°47'12.5''E); (4) Pulau Hantu (two sites: 01°13'36.5''N 103°44'47.0''E; 01°13'35.5''N 103°45'08.5''E); (5) Sisters Island (or Pulau Subar Laut; 01°12'49.0''N 103°49'59.0''E); (6) St John's Island (or Pulau Sakijang Bendera; two sites; 01°13'09.0''N 103°50'44.0''E; 01°13'25.5''N 103°50'42.5''E); (7) Lazarus Island (or Pulau Sakijang Pelepah; two sites; 01°13'12.5''N 103°51'15.0''E; 01°13'05.0''N 103°51'30.5''E) and (8) Kusu Island (or Pulau Tembakul; 01°13'33.0''N 103°51'34.0''E) (Fig. 1). Using SCUBA, meandering swim searches of approximately 60 minutes each were performed, starting from the deepest location where zooxanthellate corals could be found, usually at a depth of ~ 6 m (but ~ 9 m at Raffles Lighthouse), and ending at the shore (see DeVantier et al., 1998, 2000, 2004). Presence-absence data were collected to generate cumulative species lists of the zooxanthellate corals present at each site. Colonies that could not be identified in the field, along with new records were photographed and voucher specimens taken for identification in the laboratory. Each survey,




Fig. 1. Eight coral reefs located in the southern offshore islands of Singapore where scleractinian coral species assessments were conducted.

including the collection, generally covered an along-shore distance of 70 to 100 m. Species sightings outside of these samples but occurring in the same period were also included in the dataset.

Recent records. – Recent publications (since 2005) were analysed to extract presence-absence data for zooxanthellate coral species. Studies with relevant data were Guest et al. (2005a, 2005b), Ang (2007), Huang (2007), Huang et al. (2009) and Hoeksema (2009). With the exception of one species from Ang (2007) and Hoeksema's (2009) specimens, all records had locality information.

Museum collections and previous records. – We examined coral specimens of species deposited in the ZRC that were not found during the sampling surveys or in the work produced in the last four years. The species identities of skeletal specimens were verified and updated. For each of these species, we selected a representative specimen most identifiable based on current taxonomic descriptions as a reference for our inventory. Loo & Chou (1995) compiled a species list generated from surveys conducted during an eight-year period (1986 to 1993) and this was consulted for species not catalogued by the museum. Two species reported by Chou et al. (1994a) were also added to our list.

Species inventory. – All identifications follow Wallace (1999; see also Wallace et al., 2007) for *Acropora* spp., Hoeksema (1989) for Fungiidae, and Veron (2000) for all other zooxanthellate corals. For more detailed descriptions of corals apart from *Acropora* and Fungiidae, we referred to other taxonomic publications (Veron & Pichon, 1976, 1980, 1982; Veron et al., 1977; Veron, 1985, 1986, 1990, 2002). To obtain a comprehensive list of all species with ranges that include Singapore's reefs, we analysed the distribution maps in Wallace (1999), Hoeksema (1989) and Veron (2000). While the latter work had species ranges clearly demarcated, Wallace (1999) and Hoeksema (1989) stated the exact geographical locations where specimens were found. We set the criteria that, for reefs here to be within range of a particular *Acropora* or Fungiidae species, there must be at least two localities at approximately opposite directions of Singapore where specimens were described. Finally, we built an inventory of species in Singapore from results of the field surveys and recent records. This was supplemented by records from RMBR as well as from Chou et al. (1994a) and Loo & Chou (1995), several of which had undergone taxonomic revisions (Table 1).

RESULTS

We observed 141 species of zooxanthellate scleractinian corals during the field surveys. Including recent publications, 161 species have been encountered since 2005. Of these, 26 are newly recorded species, adding to two and five already registered in Veron (1986) and Wallace (1999), respectively. Most of the new records are from Acroporidae and Faviidae (both 33.3%). Poritidae accounts for 12.1%, Mussidae 9.1%

and Fungiidae 6.1%, while Pectiniidae and Siderastreidae each contributed 3.0% to our records. As a result, the count of coral records in Singapore reefs currently stands at 255 species from 56 genera and 15 families. This is a significant increase from the 187 zooxanthellate species listed in Loo & Chou (1995), although it still represents only 52.4% of the 487 species for which Singapore lies within the distribution ranges.

Species that were observed during the surveys and reported in recent literature since 2005 constitute 63.1% of the total number of corals ever recorded in Singapore. Of the eight reefs surveyed, Raffles Lighthouse (RL) had the highest species richness (141), followed by Pulau Jong (92), Kusu Island (91), St John's Island (74), Pulau Semakau and Pulau Hantu (both 73 species), Sisters Island (57) and Lazarus Island (55). Full data are presented in Table 2, in order of decreasing shortest straight-line distance to the mainland according to charts produced by the Hydrographic Department (2007).

DISCUSSION

The revised total number of zooxanthellate coral species recorded in Singapore is 255. This represents a 36.4% increase over the previous assessment (Loo & Chou, 1995). We also report 33 new records and present the distribution of all species encountered during our survey of eight coral reefs in Singapore (Table 2). Based on a consolidated analysis of museum records and past literature, aided by current taxonomy, we found that the inventory had 222 species prior to 2005. Thirty-five species that have been collected and deposited at the museum since the 1960s, but not recorded in the 1986–1993 transect surveys (Loo & Chou, 1995) or reported by Chou et al. (1994a), emphasize the importance of a coral reference collection. For Singapore's corals, RMBR has been an effective repository for 177 species found here. Unfortunately, newly described species such as *Goniastrea minuta* (Veron, 2000, 2002) and extremely rare species such as the endangered *Seriatopora hystrix* and *Stylophora pistillata* (see Chou et al., 1994a) may not be represented in the collection.

The distribution of new records among scleractinian families is heterogeneous. The majority of species identified for the first time in Singapore are Faviidae and Acropora corals (33.3% each). These are the result of studies by Wallace (1999), Guest et al. (2005a), Ang (2007), and Huang (2007) and Huang et al. (2009) that target specific taxonomic groups, allowing detailed evaluation of variations in coral morphology and habitat preferences. Species boundaries crucial to identification and delimitation of closely related corals can thus be clarified. Particularly in the phenotypically plastic Faviidae (e.g. Todd et al., 2001, 2002a,b, 2004), species-level assessments may omit species with overlapping morphological traits, leading to an underestimation of alpha diversity. In addition, species identification can be difficult in marginal habitats of Singapore. Corals from such areas tend to exhibit anatomical phenotypes distinct from less

impacted reefs (E. Turak & L. DeVantier, pers. comm.) where most specimen descriptions and photographs were derived (see Veron, 2000).

Changes in the recorded alpha diversity of zooxanthellate corals at three locations have occurred since the last published inventory. At Pulau Semakau, we encountered 73 zooxanthellate species in 45 genera, while 32 genera were reported in surveys from May 1986 to April 1987 (Chou & Koh, 1986; Leng et al., 1990a). At Pulau Hantu, we also observed 73 species from 45 genera, an increase over the 29 genera recorded between May 1986 and April 1987 (Chou & Koh, 1986; Leng & Lim, 1990). While 32 genera at Raffles Lighthouse were identified between September 1987 and December 1988 (Leng et al., 1990b; Goh & Chou, 1993), we found 141 species from 50 genera over the last four years. These increases are likely due to the change in survey methodology. Meandering swim searches employed in this study is a superior approach for species assessment compared to the line transect method used during the 1986–1993 surveys.

Raffles Lighthouse has the highest number of species recorded, but no clear trends exist for the other islands. Sedimentation rates increase with proximity to mainland Singapore (Lane, 1991; Low & Chou, 1994), while larval recruitment rates exhibit the opposite trend (Dikou & van Woessik, 2006), leading to the decrease in coral diversity (Chou, 1988a; Shoo, 2005; Dikou & van Woessik, 2006; Ang, 2007). Our results show that species richness is not lower for sites nearer to the mainland, although the furthest reef at Raffles Lighthouse has invariably hosted higher diversity. A greater tidal range, as compared to sites closer to the mainland, and consequently stronger water circulation at Raffles Lighthouse could depress sedimentation rates (Purchon & Enoch, 1954; Goh & Chou, 1993). Access to the reef is also restricted, making it one of the most pristine reefs in Singapore (Goh & Chou, 1993). In fact, *Stylophora pistillata*, an endangered branching species not seen since the 1960s (Chuang, 1961; Chou et al., 1994a), has been rediscovered at Raffles Lighthouse in a single stand. No data are available for entry-prohibited islands under purview of Singapore's defense ministry, but the incidental conservation may allow their reefs to serve as refuge for many marine species, including corals.

The growth form of a coral influences its response to stresses. Low light intensity in the turbid waters of Singapore may favour the foliose type of growth as the increased surface area enhances photosynthesis (Goh et al., 1994; Leng & Lim, 1990; Leng et al., 1990a; Leng et al., 1990b; Shoo, 2005). Wong (2001) found that foliose corals of the genus *Montipora* accumulate sediments faster than branching *Acropora*, but are more efficient at silt removal. Growth rates and sediment tolerance of *Acropora* are also generally lower than the other taxa (Bak & Elgershuizen, 1976; Endean, 1976; Lane, 1991; but see Rogers, 1990). Hence, as expected, our results agree with previous surveys that *Acropora* does not occur in high abundance in Singapore, and is common only at Raffles Lighthouse (Chua & Chou,

1991; Goh & Chou, 1992).

Of the 487 zooxanthellate coral species that have ranges encompassing Singapore, 255 (52.4%) have been found to be present here, either historically or recently. Although this figure is lower than that recorded in neighbouring countries, Singapore's reef area (~ 10 km²) is only approximately 0.25% of Malaysia's (4,006 km²) and 0.02% of Indonesia's (50,875 km²) reef expanses (Burke et al., 2002; Goh, 2007). Our study reveals that at least 161 species are present on the reefs here, but 94 have not been encountered recently. Land reclamation impact and associated stresses could have resulted in local extinction, but as this is not an exhaustive study—not all reefs have been surveyed—these species may still be present here. Given the large number of new species recorded in just four years, continuing survey efforts may provide further insights into the diversity and ecology of corals in Singapore.

ACKNOWLEDGEMENTS

The 14th International Marine Biology Workshop held in Singapore was organized by Tan Koh Siang (Tropical Marine Science Institute, National University of Singapore), Lena Chan (National Biodiversity Centre, National Parks Board, Singapore), Chou Loke Ming (Department of Biological Sciences, National University of Singapore) and Peter Ng (Raffles Museum of Biodiversity Research, National University of Singapore). Publication of the workshop proceedings was made possible with funds provided by the National University of Singapore and National Parks Board. We thank Wilfredo Licuanan, Emre Turak and Lyndon DeVantier for their assistance with coral identification; Bert Hoeksema and Carden Wallance for verifying our records of Fungiidae and some *Acropora* species, respectively; Kelvin Lim and Tan Siong Kiat for facilitating our access to the coral collection at RMBR; and everyone in the Marine Biology Laboratory, NUS, for their suggestions and support. We would like to dedicate this work to the late Yeo Keng Loo who had tirelessly curated the invertebrate collection and managed the coral specimens that are presented in this study.

LITERATURE CITED

- Ang, H. S., 2007. *Taxonomy and Distribution of Corals of the Family Faviidae in Singapore*. Honours thesis submitted to the Department of Biological Sciences, National University of Singapore (unpublished). 91 pp.
- Bak, R. P. M. & J. H. B. W. Elgershuizen, 1976. Patterns of oil-sediment rejection in corals. *Marine Biology*, **37**: 105–113.
- Barber, P. H., M. V. Erdmann & S. R. Palumbi, 2006. Comparative phylogeography of three codistributed stomatopods: origins and timing of regional lineage diversification in the Coral Triangle. *Evolution*, **60**: 1825–1839.
- Briggs, J. C., 1999. Coincident biogeographic patterns: Indo-West Pacific Ocean. *Evolution*, **53**: 326–335.

- Briggs, J. C., 2000. Centrifugal speciation and centres of origin. *Journal of Biogeography*, **27**: 1183–1188.
- Burke, L., E. Selig & M. Spalding, 2002. *Reefs at Risk in Southeast Asia*. World Resources Institute, Washington, DC. 72 pp.
- Carpenter, K. E. & V. G. Springer, 2005. The center of the center of marine shore fish biodiversity: the Philippine Islands. *Environmental Biology of Fishes*, **72**: 467–480.
- Chou, L. M., 1988a. Community structure of sediment stressed reefs in Singapore. *Galaxea*, **7**: 101–111.
- Chou, L. M., 1988b. The impact of human influence on the fringing reef of Pulau Hantu, Singapore. *Proceedings of the Sixth International Coral Reef Symposium, Australia*, **2**: 201–205.
- Chou, L. M., 1996. Response of Singapore reefs to land reclamation. *Galaxea*, **13**: 85–92.
- Chou, L. M., 2006. Marine habitats in one of the world's busiest harbours. In: Wolanski, E. (ed.), *The Environment in Asia Pacific Harbours*. Springer, The Netherlands. Pp. 377–391.
- Chou, L. M. & G. L. E. Koh, 1986. Initial characterisation of upper reef slope communities in Singapore waters. *Journal of the Singapore National Academy of Science*, **15**: 5–8.
- Chou, L. M., D. H. Murphy & P. K. L. Ng, 1994a. Corals, molluscs and other invertebrates. In: Ng, P. K. L. & Y. C. Wee (eds.), *The Singapore Red Data Book*. The Nature Society, Singapore. Pp. 52–87.
- Chou, L. M., J. K. Y. Low & M. G. K. Loo, 1994b. The state of coral reefs and coral reef research in Singapore. In: Wilkinson, C. R., S. Sudara & L. M. Chou (eds.), *Proceedings of the Third ASEAN-Australia Symposium on Living Coastal Resources, 16–20 May 1994, Chulalongkorn University, Bangkok, Thailand. Volume 1: Status Reviews*. Australian Institute of Marine Science, Townsville. Pp. 77–88.
- Chou, L. M., P. K. L. Ng & K. K. P. Lim, 1994c. Animalia. In: Wee, Y. C. & P. K. L. Ng (eds.), *A First Look at Biodiversity in Singapore*. National Council on the Environment, Singapore. Pp. 70–106.
- Chua, C. Y. Y. & L. M. Chou, 1991. The scleractinian community of southern islands' reefs, Singapore. In: Alcala, A. C. (ed.), *Proceedings of the Regional Symposium on Living Resources in Coastal Areas*. University of the Philippines, Manila. Pp. 41–46.
- Chuang, S. H., 1961. *On Malayan Shores*. Muwu Shosa, Singapore. 343 pp.
- Dartnall, A. & M. Jones, 1986. *A Manual of Survey Methods: Living Resources in Coastal Areas*. Australian Institute of Marine Science, Townsville. 167 pp.
- DeVantier, L. M., G. De'ath, T. J. Done & E. Turak, 1998. Ecological assessment of a complex natural system: a case study from the Great Barrier Reef. *Ecological Applications*, **8**: 480–496.
- DeVantier, L., E. Turak, K. Al-Shaikh & G. De'ath, 2000. Coral communities of the central-northern Saudi Arabian Red Sea. *Fauna of Arabia*, **18**: 23–66.
- DeVantier, L., G. De'ath, R. Klaus, S. Al-Moghrabi, M. Abdulaziz, G. B. Reinicke & C. Cheung, 2004. Reef-building corals and coral communities of the Socotra Archipelago, a zoogeographic 'crossroads' in the Arabian Sea. *Fauna of Arabia*, **20**: 117–168.
- Dikou, A. & R. van Woesik, 2006. Survival under chronic stress from sediment load: spatial patterns of hard coral communities in the southern islands of Singapore. *Marine Pollution Bulletin*, **52**: 7–21.
- Ekman, S., 1953. *Zoogeography of the Sea*. Sidgwick and Jackson, London. 417 pp.
- Endean, R., 1976. Destruction and recovery of coral reef communities. In: Jones, O. A. & R. Endean (eds.), *Biology and Geology of Coral Reefs, Volume III: Biology 2*. Academic Press, New York. Pp. 215–254.
- English, S., C. R. Wilkinson & V. Baker, 1994. *Survey Manual for Tropical Marine Resources*. Australian Institute of Marine Science, Townsville. 368 pp.
- Gilmour, J. P., 2002. Substantial asexual recruitment of mushroom corals contributes little to population genetics of adults in conditions of chronic sedimentation. *Marine Ecology Progress Series*, **235**: 81–91.
- Goh, E., 2007. *Singapore's Coral Reefs: The Next Step in its Rehabilitation and Long Term Management*. M.Sc. thesis submitted to the School of Design and Environment, National University of Singapore (unpublished). 35 pp.
- Goh, N. K. C. & L. M. Chou, 1992. A comparison of benthic life-form characteristics of a reef (Cyrene) nearest to and a reef (Raffles Lighthouse) furthest from mainland Singapore. In: Chou, L. M. & C. R. Wilkinson (eds.), *Proceedings of the Third ASEAN Science and Technology Week Conference, Vol. 6, Marine Science: Living Coastal Resources, 21–23 September, 1992, Singapore*. National University of Singapore and National Science and Technology Board, Singapore. Pp. 55–62.
- Goh, N. K. C. & L. M. Chou, 1993. The coral reef community of Pulau Satumu (Raffles Lighthouse), Singapore, with emphasis on the hard corals. *Journal of the Singapore National Academy of Science*, **20** & **21**: 51–57.
- Goh, N. K. C., C. Y. Y. Chua & L. M. Chou, 1994. Depth-related morphology of scleractinian corals on Singapore reefs. In: Sudara, S., C. R. Wilkinson & L. M. Chou (eds.), *Proceedings of the Third ASEAN-Australia Symposium on Living Coastal Resources, 16–20 May 1994, Chulalongkorn University, Bangkok, Thailand. Volume 2: Research Papers*. Australian Institute of Marine Science, Townsville. Pp. 61–67.
- Guest, J. R., A. H. Baird, B. P. L. Goh & L. M. Chou, 2005a. Reproductive seasonality in an equatorial assemblage of scleractinian corals. *Coral Reefs*, **24**: 112–116.
- Guest, J. R., A. H. Baird, B. P. L. Goh & L. M. Chou, 2005b. Seasonal reproduction in equatorial reef corals. *Invertebrate Reproduction and Development*, **48**: 207–218.
- Hoeksema, B. W., 1989. Taxonomy, phylogeny and biogeography of mushroom corals (Scleractinia: Fungiidae). *Zoologische Verhandelingen Leiden*, **254**: 1–295.
- Hoeksema, B. W., 2009. Attached mushroom corals (Scleractinia: Fungiidae) in sediment-stressed conditions at Singapore, including a new species and a new records. *Raffles Bulletin of Zoology*, Supplement No. **22**: 81–90.
- Huang, D., 2007. *Molecular and Morphological Phylogenetics of the Faviidae (Scleractinia) in Singapore*. M.Sc. thesis submitted to the Department of Biological Sciences, National University of Singapore (unpublished). 140pp.
- Huang, D., R. Meier, P. A. Todd & L. M. Chou, 2009. More evidence for pervasive paraphyly in scleractinian corals: systematic study of Southeast Asian Faviidae (Cnidaria; Scleractinia) based on molecular and morphological data. *Molecular Phylogenetics & Evolution*, **50**: 102–116.

- Koh, E. G. L. & L. M. Chou, 1989. *The Mushroom Corals of Singapore*. Department of Zoology, National University of Singapore, Singapore. 45 pp.
- Lane, D. J. W., 1991. Growth of scleractinian corals on sediment-stressed reefs at Singapore. In: Alcala, A. C. (ed.), *Proceedings of the Regional Symposium on Living Resources in Coastal Areas*. University of the Philippines, Manila. Pp. 97–106.
- Leng, C. B. & G. S. Y. Lim, 1990. An assessment of a patch reef community west of Pulau Hantu. In: Chou, L. M. (ed.), *Coastal Living Resources of Singapore: Proceedings of a Symposium on the Assessment of Living Resources in the Coastal Areas of Singapore, 3 April 1989, Singapore*. National University of Singapore, Singapore. Pp. 117–124.
- Leng, C. B., J. K. Y. Low & M. G. K. Loo, 1990a. A survey of the fringing reef community of Pulau Semakau. In: Chou, L. M. (ed.), *Coastal Living Resources of Singapore: Proceedings of a Symposium on the Assessment of Living Resources in the Coastal Areas of Singapore, 3 April 1989, Singapore*. National University of Singapore, Singapore. Pp. 101–108.
- Leng, C. B., K. S. Lee & E. G. L. Koh, 1990b. A quantitative study of the reef community at Pulau Satumu (Raffles Lighthouse). In: Chou, L. M. (ed.), *Coastal Living Resources of Singapore: Proceedings of a Symposium on the Assessment of Living Resources in the Coastal Areas of Singapore, 3 April 1989, Singapore*. National University of Singapore, Singapore. Pp. 109–116.
- Leow, W. F. L., 1992. *Taxonomy and Distribution of Acropora (Scleractinia, Acroporidae) in Singapore*. Honours project thesis submitted to the Department of Zoology, National University of Singapore (unpublished). 115 pp.
- Loo, M. G. K. & L. M. Chou, 1995. *Environmental Status of Singapore's Coral Reefs*. UNEP EAS-35 Workshop on Integrated Management of Watersheds in Relation to the Coastal and Marine Environment, 3–7 April 1995, Bangkok, Thailand. 20 pp.
- Low, J. K. Y. & L. M. Chou, 1994. Sedimentation rates in Singapore waters. In: Sudara, S., C. R. Wilkinson & L. M. Chou (eds.), *Proceedings of the Third ASEAN-Australia Symposium on Living Coastal Resources, 16–20 May 1994, Chulalongkorn University, Bangkok, Thailand. Volume 2: Research Papers*. Australian Institute of Marine Science, Townsville. Pp. 697–701.
- Hydrographic Department, 2007. *Charts for Small Craft: Singapore Strait and Adjacent Waterways*. Maritime and Port Authority, Singapore. 27 pp.
- Purchon, R. D. & I. Enoch, 1954. Zonation of the marine fauna and flora on a rocky shore near Singapore. *The Bulletin of the Raffles Museum*, **25**: 47–65.
- Rogers, C. S., 1990. Responses of coral reefs and reef organisms to sedimentation. *Marine Ecology Progress Series*, **62**: 185–202.
- Roy, K. J. & S. V. Smith, 1971. Sedimentation and coral reef development in turbid water: Fanning Lagoon. *Pacific Science*, **25**: 234–248.
- Shoo, J. H. H., 2005. *Investigating the Biodiversity of Coral Reefs in the Southern Islands of Singapore*. Honours thesis submitted to the Department of Biological Sciences, National University of Singapore (unpublished). 144 pp.
- Spalding, M. D., C. Ravilious & E. P. Green, 2001. *World Atlas of Coral Reefs*. University of California Press, California. 424 pp.
- Titlyanov, E. A. & Y. Y. Latypov, 1991. Light-dependence in scleractinian distribution in the sublittoral zone of South China Sea Islands. *Coral Reefs*, **10**: 133–138.
- Todd, P. A., P. G. Sanderson & L. M. Chou, 2001. Morphological variation in the polyps of the scleractinian coral *Favia speciosa* (Dana) around Singapore. *Hydrobiologia*, **444**: 227–235.
- Todd, P. A., R. C. Sidle & L. M. Chou, 2002a. Plastic corals from Singapore 1. *Coral Reefs*, **21**: 391–392.
- Todd, P. A., R. C. Sidle & L. M. Chou, 2002b. Plastic corals from Singapore 2. *Coral Reefs*, **21**: 407–408.
- Todd, P. A., R. J. Ladle, N. J. I. Lewin-Koh & L. M. Chou, 2004. Genotype × environment interactions in transplanted clones of the massive corals *Favia speciosa* and *Diploastrea heliophora*. *Marine Ecology Progress Series*, **271**: 167–182.
- Veron, J. E. N., 1985. New Scleractinia from Australian coral reefs. *Records of the Western Australian Museum*, **12**: 147–183.
- Veron, J. E. N., 1986. *Corals of Australia and the Indo-Pacific*. Angus & Robertson, North Ryde, Australia. 644 pp.
- Veron, J. E. N., 1990. New Scleractinia from Japan and other Indo-West Pacific countries. *Galaxea*, **9**: 95–173.
- Veron, J. E. N., 2000. *Corals of the World*. Australian Institute of Marine Science, Townsville.
- Veron, J. E. N., 2002. *New Species Described in Corals of the World*. Australian Institute of Marine Science, Townsville. 209 pp.
- Veron, J. E. N. & M. Pichon, 1976. *Scleractinia of Eastern Australia. Part I: Families Thamnasteriidae, Astrocoeniidae, Pocilloporidae*. Australian Government Publishing Service, Australia. 208 pp.
- Veron, J. E. N. & M. Pichon, 1980. *Scleractinia of Eastern Australia. Part III: Families Agariciidae, Siderastreaeidae, Fungiidae, Oculinidae, Merulinidae, Mussidae, Pectiniidae, Caryophylliidae, Dendrophylliidae*. Australian Institute of Marine Science and Australian National University Press, Australia. 471 pp.
- Veron, J. E. N. & M. Pichon, 1982. *Scleractinia of Eastern Australia. Part IV: Family Poritidae*. Australian Institute of Marine Science and Australian National University Press, Australia. 210 pp.
- Veron, J. E. N., M. Pichon & M. Wijsman-Best, 1977. *Scleractinia of Eastern Australia. Part II: Families Faviidae, Trachyphylloidae*. Australian Government Publishing Service, Australia. 231 pp.
- Wallace, C. C., 1999. *Staghorn Corals of the World*. CSIRO Publishing, Australia. 421 pp.
- Wallace, C. C., C. A. Chen, H. Fukami & P. R. Muir (2007). Recognition of separate genera within *Acropora* based on new morphological, reproductive and genetic evidence from *Acropora togianensis*, and elevation of the subgenus *Isopora* Studer, 1878 to genus (Scleractinia: Astrocoeniidae; Acroporidae). *Coral Reefs*, **26**: 231–239.
- Wong, L. S., 2001. *Assessing the Dynamics between Sedimentation, Sediment Load and Sediment Removal by Scleractinian Corals in a Singapore Fringing Reef*. Honours thesis submitted to the Department of Biological Sciences, National University of Singapore (unpublished). 51 pp.

Table 1. Synonymised species recorded in Loo & Chou (1995) and the Zoological Reference Collection (ZRC) of the Raffles Museum of Biodiversity Research at the National University of Singapore.

Previous record	Synonymy/Change	Reference
Acroporidae		
<i>Acropora brueggemanni</i>	<i>Isopora brueggemanni</i> (Brook, 1893)	Wallace et al. (2007)
<i>Acropora danai</i>	<i>Acropora abrotanoides</i> (Lamarck, 1816)	Wallace (1999)
<i>Acropora formosa</i>	<i>Acropora muricata</i> (Linnaeus, 1758)	Wallace (1999)
<i>Acropora nobilis</i>	<i>Acropora intermedia</i> (Brook, 1891)	Wallace (1999)
<i>Acropora palifera</i>	<i>Isopora palifera</i> (Lamarck, 1816)	Wallace et al. (2007)
Faviidae		
<i>Favia amicum</i>	<i>Barabattoia amicum</i> (Milne Edwards & Haime, 1850)	Veron (1986)
<i>Hydnophora exesa</i>	Family Merulinidae Verrill, 1866	Veron (1986)
<i>Hydnophora rigida</i>	Family Merulinidae Verrill, 1866	Veron (1986)
Fungiidae		
<i>Cycloseris marginata</i>	<i>Fungia (Cycloseris) costulata</i> Ortmann, 1889	Hoeksema (1989)
<i>Fungia corona</i>	<i>Fungia (Danafungia) scruposa</i> Klunzinger, 1879	Hoeksema (1989)
<i>Fungia danai</i>	<i>Fungia (Danafungia) scruposa</i> Klunzinger, 1879	Hoeksema (1989)
<i>Fungia echinata</i>	<i>Ctenactis echinata</i> (Pallas, 1776)	Hoeksema (1989)
<i>Fungia klunzingeri</i>	<i>Fungia (Danafungia) horrida</i> Dana, 1846	Hoeksema (1989)
<i>Herpetoglossa simplex</i>	<i>Ctenactis crassa</i> (Dana, 1846)	Hoeksema (1989)
<i>Lithophyllon edwardsi</i>	<i>Lithophyllon undulatum</i> Rehberg, 1892	Hoeksema (1989)

Table 2. Zooxanthellate scleractinian corals recorded in Singapore. The distribution among eight reefs of species observed during the 2006–2007 field surveys and those recorded in publications between 2005 and 2008 are shown. The source of each pre-2005 record is shown as the catalogue number and collection site, if available, for ZRC specimens, while those without specimens are referenced as Chou et al. (1994a) or Loo & Chou (1995). Localities are arranged according to decreasing shortest straight-line distance to the mainland. Abbreviations used: PR, previous record (pre-2005); NR, new record; RL, Raffles Lighthouse; SE, Pulau Semakau; JO, Pulau Jong; SI, Sisters Island; HA, Pulau Hantu; LA, Lazarus Island; SJ, St. John's Island; KU, Kusu Island; CY, Cyrene Reefs.

Species	PR	NR	Distribution (2005–2008)								Source
			RL	SE	JO	SI	HA	LA	SJ	KU	
Acroporidae											
<i>Acropora abrotanoides</i> (Lamarck, 1816)		X									Loo & Chou (1995)
<i>Acropora aculeus</i> (Dana, 1846)		X		X		X				X	
<i>Acropora acuminata</i> (Verrill, 1864)		X									ZRC.1991.600 (RL)
<i>Acropora anthocercis</i> (Brook, 1893)		X									Loo & Chou (1995)
<i>Acropora aspera</i> (Dana, 1846)			X								Wallace (1999)
<i>Acropora austera</i> (Dana, 1846)		X		X							Guest et al. (2005a)
<i>Acropora cerealis</i> (Dana, 1846)		X		X							Guest et al. (2005a)
<i>Acropora cytherea</i> (Dana, 1846)		X		X	X						
<i>Acropora dendrum</i> (Bassett-Smith, 1890)		X									ZRC.1991.597
<i>Acropora digitifera</i> (Dana, 1846)		X		X			X			X	
<i>Acropora divaricata</i> (Dana, 1846)		X									ZRC.1987.1507 (JO)
<i>Acropora donei</i> Veron & Wallace, 1984		X		X							Guest et al. (2005a)
<i>Acropora elseyi</i> (Brook, 1892)		X									ZRC.1991.611 (RL)
<i>Acropora florida</i> (Dana, 1846)			X	X							Guest et al. (2005a)
<i>Acropora glauca</i> (Brook, 1893)			X								Wallace (1999)
<i>Acropora grandis</i> (Brook, 1892)		X		X							Guest et al. (2005a)
<i>Acropora granulosa</i> (Milne Edwards & Haime, 1860)			X	X							Guest et al. (2005a)
<i>Acropora horrida</i> (Dana, 1846)		X									ZRC.1991.614 (LA)
<i>Acropora humilis</i> (Dana, 1846)		X		X							

Table 2. (Continued).

Species	PR	NR	Distribution (2005–2008)								Source
			RL	SE	JO	SI	HA	LA	SJ	KU	
<i>Acropora hyacinthus</i> (Dana, 1846)	X		X				X				Guest et al. (2005a)
<i>Acropora intermedia</i> (Brook, 1891)	X		X								
<i>Acropora latistella</i> (Brook, 1892)	X				X						
<i>Acropora longicyathus</i> (Milne Edwards & Haime, 1860)	X										ZRC.1991.624
<i>Acropora loripes</i> (Brook, 1892)		X	X								Guest et al. (2005a)
<i>Acropora lutkeni</i> Crossland, 1952	X										ZRC.1987.1510
<i>Acropora microclados</i> (Ehrenberg, 1834)		X	X								Guest et al. (2005a)
<i>Acropora microphthalma</i> (Verrill, 1869)	X										ZRC.1991.619 (RL)
<i>Acropora millepora</i> (Ehrenberg, 1834)	X		X								Loo & Chou (1995)
<i>Acropora muricata</i> (Linnaeus, 1758)	X		X								
<i>Acropora nana</i> (Studer, 1878)	X										
<i>Acropora nasuta</i> (Dana, 1846)	X		X								ZRC.1987.1511 ZRC.1987.1512 ZRC.1987.1514
<i>Acropora palmerae</i> Wells, 1954	X										
<i>Acropora polystoma</i> (Brook, 1891)	X										
<i>Acropora pulchra</i> (Brook, 1891)	X										ZRC.1992.3387 (SI)
<i>Acropora robusta</i> (Dana, 1846)	X										Wallace (1999)
<i>Acropora samoensis</i> (Brook, 1892)		X	X	X	X					X	
<i>Acropora secale</i> (Studer, 1878)	X		X		X						
<i>Acropora selago</i> (Studer, 1878)	X		X								Guest et al. (2005a)
<i>Acropora solitaryensis</i> Veron & Wallace, 1984		X									Wallace (1999)
<i>Acropora spicifera</i> (Dana, 1846)		X									Wallace (1999)
<i>Acropora subglabra</i> (Brook, 1891)		X									Veron (1986)
<i>Acropora subulata</i> (Dana, 1846)	X										ZRC.1987.2030 (HA)
<i>Acropora tenuis</i> (Dana, 1846)	X										ZRC.1991.588
<i>Acropora valenciennesi</i> (Milne Edwards & Haime, 1860)		X									Wallace (1999)
<i>Acropora valida</i> (Dana, 1846)	X		X		X	X				X	Guest et al. (2005a) ZRC.1987.2033 (JO)
<i>Acropora verweyi</i> Veron & Wallace, 1984	X		X								
<i>Acropora willisae</i> Veron & Wallace, 1984	X										
<i>Astreopora cucullata</i> Lamberts, 1980	X										Loo & Chou (1995)
<i>Astreopora expansa</i> Brüggemann, 1877	X										ZRC.1991.692 (HA)
<i>Astreopora gracilis</i> Bernard, 1896	X		X	X	X	X	X	X	X	X	ZRC.1991.691 (CY) ZRC.1991.695 (RL)
<i>Astreopora incrustans</i> Bernard, 1896	X										
<i>Astreopora listeri</i> Bernard, 1896	X										
<i>Astreopora myriophthalma</i> (Lamarck, 1816)	X		X	X	X				X	X	ZRC.1991.609 (RL) ZRC.1991.596
<i>Isopora brueggemanni</i> (Brook, 1893)	X										
<i>Isopora palifera</i> (Lamarck, 1816)	X										
<i>Montipora aequituberculata</i> Bernard, 1897	X		X		X			X	X		ZRC.1987.1521 (HA) Loo & Chou (1995)
<i>Montipora angulata</i> (Lamarck, 1816)	X										
<i>Montipora capricornis</i> Veron, 1985	X										
<i>Montipora corbettensis</i> Veron & Wallace, 1984	X		X							X	ZRC.1987.1596 (HA)
<i>Montipora crassituberculata</i> Bernard, 1897	X										
<i>Montipora danae</i> (Milne Edwards & Haime, 1851)	X		X		X	X				X	
<i>Montipora digitata</i> (Dana, 1846)	X									X	Loo & Chou (1995) Loo & Chou (1995)
<i>Montipora efflorescens</i> Bernard, 1897	X										
<i>Montipora foliosa</i> (Pallas, 1766)	X										
<i>Montipora grisea</i> Bernard, 1897	X		X	X	X	X		X		X	ZRC.1987.1600 (HA) ZRC.1987.1533 (HA)
<i>Montipora hispida</i> (Dana, 1846)	X										
<i>Montipora hoffmeisteri</i> Wells, 1954	X										
<i>Montipora informis</i> Bernard, 1897	X		X	X	X	X	X	X	X	X	ZRC.1987.1554 (HA)
<i>Montipora mollis</i> Bernard, 1897	X		X								
<i>Montipora monasteriata</i> (Forskål, 1775)	X		X	X	X	X	X			X	
<i>Montipora peltiformis</i> Bernard, 1897	X										ZRC.1987.1554 (HA)
<i>Montipora spongodes</i> Bernard, 1897	X		X		X					X	

Species	PR	NR	Distribution (2005–2008)										Source
			RL	SE	JO	SI	HA	LA	SJ	KU			
<i>Montipora spumosa</i> (Lamarck, 1816)	X								X	X			
<i>Montipora stellata</i> Bernard, 1897	X		X	X	X	X	X	X	X	X			
<i>Montipora tuberculosa</i> (Lamarck, 1816)	X		X				X						
<i>Montipora turgescens</i> Bernard, 1897	X		X	X									
<i>Montipora undata</i> Bernard, 1897	X										ZRC.1987.1533 (HA)		
<i>Montipora venosa</i> (Ehrenberg, 1834)	X										Loo & Chou (1995)		
<i>Montipora verrucosa</i> (Lamarck, 1816)	X										Loo & Chou (1995)		
Agariciidae													
<i>Gardineroseris planulata</i> (Dana, 1846)	X										Loo & Chou (1995)		
<i>Leptoseris explanata</i> Yabe & Sugiyama, 1941	X		X		X								
<i>Leptoseris hawaiiensis</i> Vaughan, 1907	X										ZRC.1987.1732 (HA)		
<i>Leptoseris scabra</i> Vaughan, 1907	X		X				X		X	X			
<i>Pachyseris rugosa</i> (Lamarck, 1801)	X		X		X								
<i>Pachyseris speciosa</i> (Dana, 1846)	X		X	X	X	X	X		X	X			
<i>Pavona cactus</i> (Forskål, 1775)	X		X										
<i>Pavona clavus</i> (Dana, 1846)	X										ZRC.1987.2094 (HA)		
<i>Pavona decussata</i> (Dana, 1846)	X		X		X	X			X	X			
<i>Pavona explanulata</i> (Lamarck, 1816)	X				X		X			X			
<i>Pavona frondifera</i> (Lamarck, 1816)	X		X	X	X		X						
<i>Pavona varians</i> Verrill, 1864	X		X										
Astrocoeniidae													
<i>Madracis kirbyi</i> Veron & Pichon, 1976	X										ZRC.1991.799 (RL)		
<i>Stylocoeniella armata</i> (Ehrenberg, 1834)	X		X	X	X		X	X	X	X			
Dendrophylliidae													
<i>Turbinaria frondens</i> (Dana, 1846)	X										Loo & Chou (1995)		
<i>Turbinaria irregularis</i> Bernard, 1896	X										Loo & Chou (1995)		
<i>Turbinaria mesenterina</i> (Lamarck, 1816)	X		X	X	X			X		X			
<i>Turbinaria peltata</i> (Esper, 1794)	X		X	X	X	X	X	X	X	X			
<i>Turbinaria radicalis</i> Bernard, 1896	X										Loo & Chou (1995)		
<i>Turbinaria reniformis</i> Bernard, 1896	X										ZRC.1991.704 (RL)		
<i>Turbinaria stellulata</i> (Lamarck, 1816)	X										ZRC.1987.2174 (HA)		
Euphyllidae													
<i>Euphyllia ancora</i> Veron & Pichon, 1980	X		X		X		X			X			
<i>Euphyllia divisa</i> Veron & Pichon, 1980	X		X										
<i>Euphyllia glabrescens</i> (Chamisso & Eysenhardt, 1821)	X										ZRC.1987.2164 (HA)		
<i>Physogyra lichtensteini</i> (Milne Edwards & Haime, 1851)	X										ZRC.1987.1889 (CY)		
<i>Plerogyra sinuosa</i> (Dana, 1846)	X		X	X	X	X	X		X	X			
Faviidae													
<i>Barabattoia amicornum</i> (Milne Edwards & Haime, 1850)	X		X		X				X				
<i>Caulastrea echinulata</i> (Milne Edwards & Haime, 1849)	X		X	X					X	X			
<i>Caulastrea furcata</i> Dana, 1846		X		X	X		X			X	Ang (2007)		
<i>Cyphastrea chalcidicum</i> (Forskål, 1775)	X		X				X	X	X				
<i>Cyphastrea microphthalma</i> (Lamarck, 1816)	X				X	X	X	X	X				
<i>Cyphastrea serailia</i> (Forskål, 1775)	X		X	X	X	X	X	X	X	X			
<i>Diploastrea heliopora</i> (Lamarck, 1816)	X		X	X	X	X	X		X	X			
<i>Echinopora gemmacea</i> Lamarck, 1816	X		X		X				X				
<i>Echinopora hirsutissima</i> Milne Edwards & Haime, 1849	X		X										
<i>Echinopora horrida</i> Dana, 1846	X										ZRC.1980.3.20.42		

Table 2 (continued).

Species	PR	NR	Distribution (2005–2008)								Source
			RL	SE	JO	SI	HA	LA	SJ	KU	
<i>Echinopora lamellosa</i> (Esper, 1795)	X		X	X	X	X	X	X	X	X	
<i>Echinopora pacificus</i> Veron, 1990		X	X		X			X		X	ZRC.CNI.0194 (RL)
<i>Favia danae</i> Verrill, 1872		X	X	X				X			ZRC.CNI.0190 (LA)
<i>Favia favius</i> (Forskål, 1775)	X		X	X	X	X	X	X	X	X	
<i>Favia laxa</i> (Klunzinger, 1879)	X		X		X		X		X	X	
<i>Favia lizardensis</i> Veron & Pichon, 1977	X		X		X			X	X	X	
<i>Favia maritima</i> (Nemenzo, 1971)	X										Loo & Chou (1995)
<i>Favia matthaii</i> Vaughan, 1918	X		X		X	X	X	X	X	X	
<i>Favia maxima</i> Veron & Pichon, 1977		X	X			X	X	X		X	ZRC.CNI.0142 (RL)
<i>Favia pallida</i> (Dana, 1846)	X		X	X	X	X			X	X	
<i>Favia rotumana</i> (Gardiner, 1899)		X	X				X		X		ZRC.CNI.0197 (RL)
<i>Favia rotundata</i> (Veron & Pichon, 1977)	X		X								Ang (2007)
<i>Favia speciosa</i> Dana, 1846	X		X	X	X	X	X	X	X	X	
<i>Favia stelligera</i> (Dana, 1846)	X				X		X				Ang (2007)
<i>Favia veroni</i> Moll & Borel-Best, 1984	X		X								Ang (2007)
<i>Favites abdita</i> (Ellis & Solander, 1786)	X		X	X	X	X	X	X	X	X	
<i>Favites chinensis</i> (Verrill, 1866)	X		X				X	X	X	X	
<i>Favites complanata</i> (Ehrenberg, 1834)	X		X		X	X		X	X	X	
<i>Favites flexuosa</i> (Dana, 1846)	X		X		X			X	X		
<i>Favites halicora</i> (Ehrenberg, 1834)	X		X		X	X	X	X	X	X	
<i>Favites paraflexuosa</i> Veron, 2000		X				X		X	X		ZRC.CNI.0165 (SI)
<i>Favites pentagona</i> (Esper, 1794)	X		X				X	X	X	X	
<i>Favites russelli</i> (Wells, 1954)	X										Loo & Chou (1995)
<i>Goniastrea aspera</i> Verrill, 1905	X		X	X	X	X	X			X	
<i>Goniastrea australensis</i> (Milne Edwards & Haime, 1857)	X		X		X	X				X	
<i>Goniastrea edwardsi</i> Chevalier, 1971	X		X	X	X	X		X		X	
<i>Goniastrea favulus</i> (Dana, 1846)	X		X					X	X		
<i>Goniastrea minuta</i> Veron, 2000		X	X				X				Guest et al. (2005b)
<i>Goniastrea palauensis</i> (Yabe & Sugiyama, 1936)	X		X						X		
<i>Goniastrea pectinata</i> (Ehrenberg, 1834)	X		X	X	X	X	X	X	X	X	
<i>Goniastrea retiformis</i> (Lamarck, 1816)	X		X	X	X	X		X	X	X	
<i>Leptastrea pruinosa</i> Crossland, 1952	X		X	X				X		X	
<i>Leptastrea purpurea</i> (Dana, 1846)	X		X	X	X	X	X	X	X		
<i>Leptastrea transversa</i> Klunzinger, 1879	X		X	X	X	X	X		X	X	
<i>Leptoria phrygia</i> (Ellis & Solander, 1786)	X		X		X					X	
<i>Montastrea annuligera</i> (Milne Edwards & Haime, 1849)	X										Loo & Chou (1995)
<i>Montastrea colemani</i> Veron, 2000		X	X								ZRC.CNI.0237 (RL)
<i>Montastrea curta</i> (Dana, 1846)	X		X			X	X		X	X	
<i>Montastrea magnistellata</i> Chevalier, 1971	X		X		X	X	X	X	X	X	
<i>Montastrea valenciennesi</i> (Milne Edwards & Haime, 1848)	X		X	X	X	X	X	X	X	X	
<i>Oulastrea crispata</i> (Lamarck, 1816)	X		X	X	X		X	X		X	
<i>Oulophyllia bennettiae</i> (Veron & Pichon, 1977)		X	X	X		X			X	X	ZRC.CNI.0169 (SJ)
<i>Oulophyllia crispa</i> (Lamarck, 1816)		X	X			X				X	ZRC.CNI.0145 (RL)
<i>Platygyra daedalea</i> (Ellis & Solander, 1786)	X		X	X	X	X		X	X	X	
<i>Platygyra lamellina</i> (Ehrenberg, 1834)	X		X						X		
<i>Platygyra pini</i> Chevalier, 1975	X		X	X	X	X	X		X	X	
<i>Platygyra ryukyuensis</i> Yabe & Sugiyama, 1936		X	X				X				Guest et al. (2005b)
<i>Platygyra sinensis</i> (Edwards & Haime, 1849)	X		X	X	X	X	X	X	X	X	
<i>Platygyra verweyi</i> Wijsman-Best, 1976	X		X		X	X			X	X	
Fungiidae											
<i>Ctenactis albitentaculata</i> Hoeksema, 1989	X										ZRC.1987.1749
<i>Ctenactis crassa</i> (Dana, 1846)	X										ZRC.1987.3578 (HA)

Table 2 (continued).

Species	PR	NR	Distribution (2005–2008)								Source
			RL	SE	JO	SI	HA	LA	SJ	KU	
<i>Ctenactis echinata</i> (Pallas, 1766)	X		X	X	X		X			X	Loo & Chou (1995) ZRC.1987.3493 (HA)
<i>Fungia (Cycloseris) vaughani</i> Boschma, 1923	X										
<i>Fungia (Danafungia) horrida</i> Dana, 1846	X										
<i>Fungia (Danafungia) scruposa</i> Klunzinger, 1879	X		X	X	X	X					Loo & Chou (1995) Hoeksema (2009)
<i>Fungia (Fungia) fungites</i> (Linnaeus, 1758)	X		X		X	X	X		X	X	
<i>Fungia (Lobactis) scutaria</i> Lamarck, 1801	X										
<i>Fungia (Pleuractis) moluccensis</i> van der Horst, 1919	X										Loo & Chou (1995) Hoeksema (2009)
<i>Fungia (Pleuractis) paumotensis</i> Stutchbury, 1833	X		X	X	X	X		X	X	X	
<i>Fungia (Verrillofungia) concinna</i> Verrill, 1864	X			X		X		X	X	X	
<i>Fungia (Verrillofungia) repanda</i> Dana, 1846	X		X	X	X		X	X	X	X	ZRC.1987.2206 (CY)
<i>Fungia (Verrillofungia) scabra</i> Döderlein, 1901	X				X		X		X	X	
<i>Fungia (Wellsofungia) granulosa</i> Klunzinger, 1879	X										
<i>Heliofungia actiniformis</i> (Quoy & Gaimard, 1833)	X		X	X	X		X				Hoeksema (2009)
<i>Herpolitha limax</i> (Esper, 1797)	X		X	X	X		X		X		
<i>Lithophyllon undulatum</i> Rehberg, 1892	X		X	X	X		X	X	X	X	
<i>Podabacia crustacea</i> (Pallas, 1766)	X		X	X	X	X	X		X	X	Hoeksema (2009)
<i>Podabacia kunzmanni</i> Hoeksema, 2009		X									
<i>Podabacia motuporensis</i> Veron, 1990		X									
<i>Polyphyllia talpina</i> (Lamarck, 1801)	X			X			X				
Merulinidae											
<i>Hydnophora exesa</i> (Pallas, 1766)	X		X	X	X		X	X		X	ZRC.CNI.0238 (SE) ZRC.1987.1829 (JO) Loo & Chou (1995)
<i>Hydnophora rigida</i> (Dana, 1846)	X		X							X	
<i>Merulina ampliata</i> (Ellis & Solander, 1786)	X		X	X	X	X	X	X	X	X	
<i>Merulina scabricula</i> Dana, 1846	X		X						X	X	ZRC.CNI.0239 (SJ) Veron (1986)
<i>Scapophyllia cylindrica</i> Milne Edwards & Haime, 1848	X		X							X	
Mussidae											
<i>Acanthastrea echinata</i> (Dana, 1846)	X			X	X		X		X	X	ZRC.CNI.0238 (SE) ZRC.1987.1829 (JO) Loo & Chou (1995)
<i>Acanthastrea hillae</i> Wells, 1955	X		X				X	X		X	
<i>Acanthastrea rotundoflora</i> Chevalier, 1975		X		X	X						
<i>Lobophyllia corymbosa</i> (Forskål, 1775)	X										ZRC.CNI.0239 (SJ) Veron (1986)
<i>Lobophyllia hataii</i> Yabe & Sugiyama, 1936	X										
<i>Lobophyllia hemprichii</i> (Ehrenberg, 1834)	X		X	X	X		X		X	X	
<i>Symphyllia agaricia</i> Milne Edwards & Haime, 1849		X	X						X		ZRC.CNI.0239 (SJ) Veron (1986)
<i>Symphyllia radians</i> Milne Edwards & Haime, 1849	X		X	X	X	X	X	X	X	X	
<i>Symphyllia recta</i> (Dana, 1846)	X		X	X	X		X		X	X	
<i>Symphyllia valenciennesii</i> Milne Edwards & Haime, 1849		X									Veron (1986)
Oculinidae											
<i>Galaxea astreata</i> (Lamarck, 1816)	X		X								Guest et al. (2005b)
<i>Galaxea fascicularis</i> (Linnaeus, 1767)	X		X		X	X	X	X	X	X	
Pectiniidae											
<i>Echinophyllia aspera</i> (Eillis & Solander, 1788)	X		X	X		X	X	X	X	X	ZRC.1987.2146 Loo & Chou (1995)
<i>Echinophyllia echinata</i> (Saville-Kent, 1871)	X										
<i>Echinophyllia echinoporoides</i> Veron & Pichon, 1980	X										
<i>Mycedium elephantotus</i> (Pallas, 1766)	X		X	X	X	X	X			X	Loo & Chou (1995) ZRC.1987.1845 (SE)
<i>Oxypora crassispinosa</i> Nemenzo, 1979	X										
<i>Oxypora glabra</i> Nemenzo, 1959	X										
<i>Oxypora lacera</i> (Verrill, 1864)	X		X	X	X		X				ZRC.CNI.0240 (SE)
<i>Pectinia alcornis</i> (Saville-Kent, 1871)	X		X	X	X	X	X		X	X	
<i>Pectinia ayleni</i> (Wells, 1935)		X		X							

Table 2 (continued).

Species	PR	NR	Distribution (2005–2008)									Source
			RL	SE	JO	SI	HA	LA	SJ	KU		
<i>Pectinia lactuca</i> (Pallas, 1766)	X		X		X							
<i>Pectinia paeonia</i> (Dana, 1846)	X		X	X	X		X					
Pocilloporidae												
<i>Pocillopora damicornis</i> (Linneaus, 1758)	X		X	X	X	X	X	X	X	X		
<i>Pocillopora verrucosa</i> (Ellis & Solander, 1786)	X										ZRC.1991.625	
<i>Seriatopora hystrix</i> Dana, 1846	X										Chou et al. (1994a)	
<i>Stylophora pistillata</i> Esper, 1797	X		X									
Poritidae												
<i>Alveopora allingi</i> Hoffmeister, 1925	X										Loo & Chou (1995)	
<i>Alveopora catalai</i> Wells, 1968	X										ZRC.1987.1890 (HA)	
<i>Alveopora excelsa</i> Verrill, 1863	X										ZRC.1980.3.20.120	
<i>Alveopora fenestrata</i> (Lamarck, 1816)	X										ZRC.1987.1892 (HA)	
<i>Alveopora marionensis</i> Veron & Pichon, 1982	X										Loo & Chou (1995)	
<i>Alveopora spongiosa</i> Dana, 1846		X	X	X							ZRC.CNI.0241 (RL)	
<i>Alveopora tizardi</i> Bassett-Smith, 1890		X	X	X			X			X	ZRC.CNI.0245 (SE)	
<i>Goniopora columna</i> Dana, 1846	X		X	X	X	X		X		X		
<i>Goniopora djiboutiensis</i> Vaughan, 1907	X										ZRC.1987.2184 (CY)	
<i>Goniopora eclipsensis</i> Veron & Pichon, 1982	X										ZRC.1987.1905 (HA)	
<i>Goniopora fruticosa</i> Saville-Kent, 1893	X										ZRC.1987.1903 (CY)	
<i>Goniopora lobata</i> Milne Edwards & Haime, 1860	X		X		X				X			
<i>Goniopora pandoraensis</i> Veron & Pichon, 1982	X										ZRC.1987.1909	
<i>Goniopora pendulus</i> Veron, 1985	X										ZRC.1987.1916 (HA)	
<i>Goniopora somaliensis</i> Vaughan, 1907	X										ZRC.1987.1914 (HA)	
<i>Goniopora stokesi</i> Milne Edwards & Haime, 1851	X										Loo & Chou (1995)	
<i>Goniopora stutchburyi</i> Wells, 1955	X										ZRC.1980.3.20.130	
<i>Porites australiensis</i> Vaughan, 1918	X			X	X			X	X			
<i>Porites cylindrica</i> Dana, 1846	X										ZRC.1987.1925 (SE)	
<i>Porites deformis</i> Nemenzo, 1955		X	X	X							ZRC.CNI.0242 (RL)	
<i>Porites lichen</i> Dana ,1846	X				X							
<i>Porites lobata</i> Dana, 1846	X		X	X	X	X	X	X	X	X		
<i>Porites lutea</i> Milne Edwards & Haime, 1851	X		X	X	X	X	X		X	X		
<i>Porites monticulosa</i> Dana, 1846		X	X			X		X	X	X	ZRC.CNI.0243 (RL)	
<i>Porites murrayensis</i> Vaughan, 1918	X										Loo & Chou (1995)	
<i>Porites nigrescens</i> Dana, 1846	X										ZRC.1987.1951 (HA)	
<i>Porites rus</i> (Forskål, 1775)	X										ZRC.1987.1961 (SE)	
<i>Porites solida</i> (Forskål, 1775)	X										ZRC.1980.3.20.132	
<i>Porites stephensoni</i> Crossland, 1952	X										ZRC.1987.1936 (HA)	
<i>Porites vaughani</i> Crossland, 1952	X										Loo & Chou (1995)	
Siderastreidae												
<i>Coscinaraea columna</i> (Dana, 1846)	X		X	X	X	X		X		X		
<i>Coscinaraea exesa</i> (Dana, 1846)	X										Loo & Chou (1995)	
<i>Psammocora contigua</i> (Esper, 1797)	X										ZRC.1989.41 (HA)	
<i>Psammocora digitata</i> Milne Edwards & Haime, 1851	X		X	X	X		X			X		
<i>Psammocora explanulata</i> van der Horst, 1922	X										Loo & Chou (1995)	
<i>Psammocora superficialis</i> Gardiner, 1898	X							X		X		
<i>Pseudosiderastrea tayami</i> Yabe & Sugiyama, 1935		X	X	X	X		X	X	X		ZRC.CNI.0244 (RL)	
Trachyphylliidae												
<i>Trachyphyllia geoffroyi</i> (Audouin, 1826)	X		X	X			X					